arXiv:gr-qc/0509042v1 13 Sep 2005

The Closed Universe Model with Variable $\Lambda-$term

P.I. Fomin[a] , P.A.Nakaznoy[b] , S.I.Vilchinskyi[b]
a - Bogolyubov Institute for Theoretical Physics, Kiev, Ukraine
b - Kiev National Taras Schevchenko University, Ukraine
pfomin@bitp.kiev.ua , nak@univ.kiev.ua , sivil@univ.kiev.ua

Abstract

We have studied the closed universe model with the variable cosmological term, which is presented as a sum of two terms: $\Lambda = \Lambda_0 - kR$. First term Λ_0 is a constant and it is describing a sum of quantum field's zero oscillations. In the geometrical sense it determines an own constant curvature of the space-time. The second term is variable, it is proportional to curvature of the space-time and can be interpretable as response of vacuum to curvature of the space-time. In the framework of such model we have investigated the time dynamic of $\Lambda-$term and try to explain a mechanism which to give rise to a large value of Λ hence leading to the "cosmological constant problem". It was studied models for different dark matter equation of state: $p = 0$ and $p = \varepsilon$. It was shown that the presence of a variable $\Lambda-$term alters the evolution of the scale factor a in the very early epoch. In the context of this model, a severe decrease of the module of vacuum energy density at the primitive stage of Universe's development is explained. The numerical results for the value of the Λ-term agree with theoretical predictions for the early epoch and with recent observation data.

1 Introduction

Recent observation of Type 1a supernova (based on high red-shift supernova) [1]-[3] and structure formation models [4], [5] indicating an accelerating universe have once more drawn attention to the possible existence of a small positive cosmological $\Lambda-$term. As well known, Λ-term can produce the field of anti-gravity that causes the cosmological expansion to accelerate (see, for example [6]). At a theoretical level a cosmological term is predicted to arise out of zero-point quantum vacuum fluctuation of fundamental scalar, spinor, vector and tensor fields. Although a theoretical predicted value of the energy density of vacuum usually appears to be much larger than current observation limit [6]. One method of resolving the dilemma between a very large cosmological term, predicted by field theory and an extremely small one, suggested observation with obvious advantages is to make the cosmological term time-depended.

At the present moment there are several models with variable Λ-term and next evolutionary relation for Λ-term [7]:

$$\Lambda \sim t^{-2}, \quad \Lambda \sim t^{-\alpha}, \quad \Lambda \sim A + B\exp(-\alpha t), \quad \Lambda \sim a^{-2}, \quad \Lambda \sim T^{\alpha}, \quad \Lambda \sim a^{-\alpha},$$

$$\Lambda \sim \exp(-\alpha a), \quad \Lambda \sim H^2, \quad \Lambda \sim H^2 + Aa^{-\alpha}, \quad \Lambda \sim f(H), \quad \frac{d\Lambda}{dt} \sim g(\Lambda, H)$$

where a is the scale factor, H is the Hubble parameter, T is the temperature, t is cosmic time, A, B, α are the constants. However the most of these models are not necessarily backed by strong physical arguments and does not lead to adequate physical results.

In this work we have studied the closed universe model with variable cosmological term which is presented as a sum of two terms. We assume that

$$\Lambda = \Lambda_0 - kR \tag{1}$$

where $\Lambda_0, \quad k$ are constants, R is scalar curvature of space-time on the basic next physical arguments. As well know Einstein's equations can be derived from the action [8]

$$A = \frac{c^4}{16\pi G}\int R\sqrt{-g}d^4x + \frac{1}{c}\int L_{mat}(\varphi, \partial_\nu\varphi)\sqrt{-g}d^4x,$$

where L_{mat} is the Lagrangian for matter depending some dynamical variables generically denoted as φ. The variation of this action with respect to these variables φ will lead to the equation of motion for matter in a given background geometry, while the variation of the action with respect to the metric tensor $g_{\alpha\beta}$ leads to the Einstein's equations:

$$R_{\alpha\beta} - \frac{1}{2}g_{\alpha\beta}R = \frac{8\pi G}{c^4}T_{\alpha\beta}.$$

As is well known (see for example [7]), the consideration a new matter action $\acute{L}_{mat} = L_{mat} - \frac{\Lambda}{\frac{8\pi G}{c^4}}$ where Λ is constant, does not lead to change of the equation of motion for the matter under this transformation, but the action will picks up an extra term proportional to Λ :

$$A = \frac{c^4}{16\pi G}\int R\sqrt{-g}d^4x + \frac{1}{c}\int (L_{mat} - \frac{\Lambda c^4}{8\pi G})\sqrt{-g}d^4x = \\ = \frac{c^4}{16\pi G}\int (R - 2\Lambda)\sqrt{-g}d^4x + \frac{1}{c}\int L_{mat}\sqrt{-g}d^4x \qquad (2)$$

and Einstein's equations gets modified – the "Λ-term"appears. The physical interpretation of λ-term depending which of the two forms of the equation (2) for the action is used. According to choice words somebody, the cosmological constant has the many faces. The first interpretation, based on the first line of equation (2), treats Λ as the shift in the matter Lagrangian or, the same, the shift in the matter Hamiltonian. Such a constant shift in the energy does not affect the dynamics of matter, while gravity, which couples to the total energy of the system, picks up an extra contribution in the form of a new term in the energy-momentum tensor, and Einstein's equations can be write in such form:

$$R_{\alpha\beta} - \frac{1}{2}g_{\alpha\beta}R = \frac{8\pi G}{c^4}(T_{\alpha\beta} + \frac{c^4}{8\pi G}\Lambda g_{\alpha\beta}).$$

In such interpretation introducing the cosmological term is like introducing a cosmological vacuum with energy momentum tensor $T^{vac}_{\alpha\beta}$ given by $T^{vac}_{\alpha\beta} = \frac{c^4}{8\pi G}\Lambda g_{\alpha\beta}$. In other words – if the vacuum has energy it also gravitates. The quantum field theory proves that vacuum has energy [9] and what is more quantum field theory in curved space-time predicted that the density energy of vacuum is proportional to scalar curvature in curved space-time [10], [11]. However, numerical calculations of the vacuum energy density in this case, with account for radiative corrections, lead to divergence even in the single-loop approximation [11]. All these results were obtained in a continuous space-time. The situation changes drastically if one recalls that, as was shown by P.Fomin [12], taking into account self-gravitation makes the physical vacuum unstable relative to the "decay"into a set of long-living coupled "Planck cells". Every "Planck cell"represents a microscopic closed universe with zero total energy. In other words, taking into account self-gravitation of vacuum fluctuations leads to a discrete (lattice) structure of the vacuum at Planck sizes. Each such configuration has zero total mass, charge, and energy, but a nonzero multipole momentum, which leads to a gravitational coupling between "Planck cells", and therefore to their curving. A curving at the level of Planck cells causes the appearance of a nonzero finite vacuum energy, proportional to the scalar curvature (the vacuum energy must be Lorentz covariant!). Hence, within such an approach one can assume that accounting for self-gravitation of vacuum fluctuations leads to a discrete (quantized) contribution to the scalar curvature and the vacuum energy density is proportional to the scalar curvature: $\Lambda \sim R$.

This approach, in an interesting way, is in accordance with an idea once expressed by A.D.Sakharov [13], who suggested to treat scalar curvature as the rigidity of space-time.

So, in this interpretation we assume that

$$T^{vac}_{\alpha\beta} = \frac{1}{\varkappa}(\Lambda_0 - kR)g_{ik} \equiv \lambda_0 - \frac{k}{\varkappa}R,$$

where $\varkappa = \dfrac{8\pi G}{c^4}$. The parameter k makes a sence of space's elastic, Λ_0 responds to the sum of quantum field's zero-fluctuations.

On the other side the second line in equation (2) can be interpreted as gravitational field, determined by the Lagrangian of the form $L_{grav} \sim \dfrac{1}{G}(R - 2\Lambda)$ and interaction with matter described by the Lagrangian L_{matter}. In this interpretation, gravity is described by two parameters - the Newton's constant G and the cosmological term Λ. It is then to modify the left hand side of Einstein's equations and write

$$R_{\alpha\beta} - \frac{1}{2}g_{\alpha\beta}R - g_{\alpha\beta}\Lambda = \frac{8\pi G}{c^4}T_{\alpha\beta}.$$

It is important that in this interpretation, the space-time is treated as curved even in the absence of matter, since the equation

$$R_{\alpha\beta} - \frac{1}{2}g_{\alpha\beta}R - g_{\alpha\beta}\Lambda = 0$$

does not admit flat space-time as solution. Therefore in this case $\Lambda-$term has the geometrical nature – it determines an own constant curvature of the space-time and it is constant – Λ_0.

2 Basic equations

In fact, it is possible to consider a situation in which both effects can occur. If the matter Lagrangian contains energy densities of vacuum, which is proportional to scalar curvature, then $\Lambda-$ term can change due to the Richi scalar R changes during the evolution. On the other side, if the gravitational interaction is described by the Lagrangian of the form $\frac{1}{G}(R-2\Lambda)$, then there is an intrinsic cosmological constant as the primordial own curvature of space-time in nature just as there is a Newtonian gravitational constant in nature. Hence, within such an approach we will review the universe model with cosmological term which is presented as a sum of two terms (1).

In framework of such assumption the Einstein's equations will be have next view:

$$R_{\alpha\beta}-\frac{1}{2}g_{\alpha\beta}R-\Lambda_0 g_{\alpha\beta}=\varkappa T_{\alpha\beta}-kRg_{\alpha\beta}\ , \tag{3}$$

where $\varkappa=\frac{8\pi\gamma}{c^4}$, $T_{\alpha\beta}=(\varepsilon+p)\,u_\alpha\,u_\beta-pg_{\alpha\beta}$ – energy-momentum tensor for a perfect fluid, ϵ is the mass-energy density of matter, p is the pressure and u^α is the four-velocity of matter.

We assume the closed homogeneous and isotropic universe and used the Friedmann-Robertson-Walker metric:

$$ds^2=c^2dt^2-a^2(t)\left(d\chi^2+\sin^2\chi\left(d\theta^2+\sin^2\theta\,d\phi^2\right)\right)\ . \tag{4}$$

In this denotations we have this Friedmann's equations:

$$\begin{cases}\dot{a}^2+c^2=\frac{1}{3}\varkappa c^2a^2(\varepsilon+\varepsilon_V)\\ \ddot{a}=-\frac{1}{6}\varkappa c^2a(\varepsilon+3p-2\varepsilon_V)\ .\end{cases} \tag{5}$$

For our model (1) from this follows:

$$\begin{cases}\dot{a}^2+fc^2=\dfrac{\varkappa c^2}{1-4k}a^2\left(\varepsilon\left(\dfrac{1}{3}-k\right)-pk+\dfrac{\lambda_0}{3}\right)\\ \ddot{a}=-\dfrac{\varkappa c^2}{1-4k}a\left(\varepsilon\left(\dfrac{1}{6}-k\right)+p\left(\dfrac{1}{2}-k\right)-\dfrac{\lambda_0}{3}\right)\ ,\\ p=\nu\varepsilon\ ,\end{cases} \tag{6}$$

where ν_i is determined the equation state selection: $\nu = 0$ for zero pressure, $\nu = \frac{1}{3}$ for pure radiation and $\nu = 1$ for limited hard state of a matter.

From Fridman's equations (6) following the principle of energy density changing

$$\varepsilon a^{\sigma} = Const, \qquad \sigma = \frac{3(1-4k)(\nu+1)}{1-3k(\nu+1)}. \tag{7}$$

According to different equation of state σ is also different, therefore:

$$\begin{cases} \text{dust: } \nu = 0 \Longrightarrow \varepsilon a^{\frac{3(1-4k)}{1-3k}} = \varepsilon_0 a_0^{\frac{3(1-4k)}{1-3k}} \\ \text{pure radiation: } \nu = \dfrac{1}{3} \Longrightarrow \varepsilon a^4 = \varepsilon_0 a_0^4 \\ \text{limited hard state of a matter: } \nu = 1 \Longrightarrow \varepsilon a^{\frac{6(1-4k)}{1-6k}} = \varepsilon_0 a_0^{\frac{6(1-4k)}{1-6k}}, \end{cases} \tag{8}$$

where ε_0 – modern values of the matter density for different kind of matter, $a_0 = \frac{c}{H_0}$ – modern value of the Hubble radius, $H_0 = 65 \pm 15 \frac{\text{km}}{\text{sec Mps}}$ – modern value of the Hubble parameter. For next evaluations we are going to consider, that

$$\varepsilon = \sum_i \varepsilon_i \ , p = \sum_i \nu_i \varepsilon_i \ .$$

First of all, lets research a concordance of our model with the data of last observations [14]:

type of matter	relative density $\Omega_{i0} = \frac{\rho_{i0}}{\rho_c}$
vacuum	$\Omega_{\Lambda 0} = 0.7 \pm 0.1$
dark matter	$\Omega_{D0} = 0.3 \pm 0.1$
baryon matter	$\Omega_{B0} = 0.02 \pm 0.01$
radiation	$\Omega_{R0} = 0.8 \cdot 10^{-4} \alpha \ , \alpha \backsim 1 \div 10$

There

$$\Omega = \frac{\rho}{\rho_c} \ , \tag{9}$$

$$\rho_c = \frac{3{H_0}^2}{8\pi G} = (0.6 \pm 0.1) \times 10^{-29} \frac{\text{g}}{\text{sm}^3} \tag{10}$$

$$H_0 = 65 \pm 15 \frac{\text{km}}{\text{s} \cdot \text{Mpk}} \text{ - Hubble's constant .} \tag{11}$$

Certainly, use this data for our model is an approximation. But, it is a good approximation, because now we have very small $R \sim \Lambda$.

Lets equate any appropriate equations of the systems (5) and (6). As a result we are taken:

$$\Omega_0 = \Omega_{\Lambda 0} - k(4\Omega_{\Lambda 0} + \Omega_{D0}(1 - 3\nu_D) + \Omega_{B0}) \ , \tag{12}$$

where ν_D is the parameter of dark matter state's equation. We use $p = 0$ for a barion matter and $p = \dfrac{\varepsilon}{3}$ for a radiation. We don't know a nature of the dark matter, therefore use for it this general state's equation.

Under $z \cong 0.7$, allegedly, we have zero acceleration of the Universe [14]. Hence we can obtain unknown k. For this equate acceleration under this z to zero, the principle of energy density changing is taking into account. As a result we are taken this transcendental equation for k:

$$k = \frac{\Omega_D(1 + 3\nu_D) + \Omega_B + 2\Omega_R - 2\Omega_{\Lambda 0}}{2(3\Omega_D(1 + \nu_D) + 3\Omega_B + 4\Omega_R - 4\Omega_{\Lambda 0} - \Omega_{D0}(1 - 3\nu_D) - \Omega_{B0})} \ , \tag{13}$$

where $\Omega_i = \Omega_{i0}(1 + z)^{\sigma_i}$.

For numerical calculations we use two possible state's equations of the dark matter: $\nu_D = 0$, i.e. dust and $\nu_D = 1$, which is fiting with limit hard state of matter [16]. For this two cases from (13) follows:

$$\begin{cases} \nu_D = 0 \Longrightarrow k = 0.03 \ , \Omega_0 = 0.6 \\ \nu_D = 1 \Longrightarrow k = 0.03 \ , \Omega_0 = 0.63 \ . \end{cases} \tag{14}$$

Comment, that we have equal values k within the bounds of modern data.

For the next calculations we use $a_0 = \dfrac{c}{H_0}$, i.e. use hubble's radius. The principle of energy density changing will be considered to be true for every kind of matter. Then we can evaluate Λ-term as function from a:

$$\Lambda = \frac{3\Omega_0}{(1 - 4k){a_0}^2} + \frac{3k}{1 - 4k}\left((1 - 3\nu_D)\Omega_{D0}\frac{{a_0}^{\sigma_D - 2}}{a^{\sigma_D}} + \Omega_{B0}\frac{{a_0}^{\sigma_B - 2}}{a^{\sigma_B}}\right) \ . \tag{15}$$



Figure 1: The dependence of the non-dimensional Λ-term $\Omega_\Lambda = \dfrac{\Lambda}{\varkappa\rho_c c^2}$ from the $x = \dfrac{a}{a_0}$ for the barion model of a dark model, $k = 0.03$, $\Omega_0 = 0.6$.

For the concerned cases:

$$\begin{cases} \nu_D = 0 \Longrightarrow \Lambda = \dfrac{3\Omega_0}{(1-4k){a_0}^2} + \dfrac{3k}{1-4k}(\Omega_D + \Omega_B)\dfrac{{a_0}^{\sigma_0 - 2}}{a^{\sigma_0}} \ , \sigma_0 = \dfrac{3(1-4k)}{1-3k} \\ \nu_D = 1 \Longrightarrow \Lambda = \dfrac{3\Omega_0}{(1-4k){a_0}^2} + \dfrac{3k}{1-4k}\left(\Omega_B\dfrac{{a_0}^{\sigma_0-2}}{a^{\sigma_0}} - 2\Omega_D\dfrac{{a_0}^{\sigma_1-2}}{a^{\sigma_1}}\right) \ , \sigma_1 = \dfrac{6(1-4k)}{1-6k} \ . \end{cases} \tag{16}$$

From (16) it follows, that Λ-term is positive, when $0 < k < \dfrac{1}{3}$ for the barion model of a dark matter and $k < 0$ for limited hard state. This is means, that limited hard state of a matter is requires with value $k = 0.03$ for a negative Λ-term in the development of Universe.

For the purposes of illustrations (16) we make next two plots, which describing the dependence of the non-dimensional Λ-term $\Omega_\Lambda = \dfrac{\Lambda}{\varkappa\rho_c c^2}$ from the $x = \dfrac{a}{a_0}$.

With the help of the principle of energy density changing (7) we can write the integral for $a(t)$. In non-dimeonsional variables $x = \dfrac{a}{a_0}$ and $t = \tau H_0$ it has such form:

$$\tau = \int_0^x \frac{dx'}{\sqrt{\dfrac{1}{1-4k}\sum_i (1 - 3k(\nu_i + 1))\Omega_{i0}{x'}^{(2-\sigma_i)} + \dfrac{\Omega_0}{1-4k}{x'}^2 - 1}} \ . \tag{17}$$



Figure 2: The dependence of the non-dimensional Λ-term $\Omega_\Lambda = \dfrac{\Lambda}{\varkappa\rho_c c^2}$ from the $x = \dfrac{a}{a_0}$ for the limited hard model of a dark model $k = 0.03$, $\Omega_0 = 0.63$.

For the two considered models of a dark matter:

$$\begin{cases} \nu_D = 0 \Longrightarrow \\ \tau = \displaystyle\int_0^x \frac{dx'}{\sqrt{\dfrac{1-3k}{1-4k}(\Omega_{D0}+\Omega_{B0})\; x'^{-\dfrac{1-6k}{1-3k}} + \dfrac{\Omega_{R0}}{x'^2} + \dfrac{\Omega_0}{1-4k}x'^2 - 1}} \\ \nu_D = 1 \Longrightarrow \\ \tau = \displaystyle\int_0^x \frac{dx'}{\sqrt{\dfrac{1-6k}{1-4k}\Omega_{D0}x'^{-\dfrac{4(1-3k)}{1-6k}} + \dfrac{1-3k}{1-4k}\Omega_{B0}\; x'^{-\dfrac{1-6k}{1-3k}} + \dfrac{\Omega_{R0}}{x'^2} + \dfrac{\Omega_0}{1-4k}x'^2 - 1}} \; . \end{cases} \tag{18}$$

The results of numerical calculations of this integrals presents at the following plot.

Under little x this integrals may be calculate in analytic form and solve to x:

$$\begin{cases} \nu_D = 0 \Longrightarrow x(\tau) = (\Omega_{R0})^{\dfrac{1}{4}}\sqrt{2\tau} \\ \nu_D = 1 \Longrightarrow x(\tau) = [\sqrt{\dfrac{1-4k}{1-6k}\Omega_{D0}}\; 3\tau]^{\dfrac{1-6k}{3(1-4k)}} \; . \end{cases} \tag{19}$$

The results of calculations are reducing to next tables. The table №1 is responding to the barion model of a dark matter ($k = 0.03$, $\Omega_0 = 0.6$) , №2 - limited hard ($k = 0.03$, $\Omega_0 = 0.63$) .



Figure 3: The dependence of the non-dimensional variable $x = \dfrac{a}{a_0}$ from the non-dimensional time $\tau = tH_0$. The lower curve is responding to the barion model of a dark matter, $k = 0.03$, $\Omega_0 = 0.6$. The high is responding to the limited hard model of a dark matter, $k = 0.03$, $\Omega_0 = 0.63$. The medium is responding to the standard cosmological model with a flat universe and constant cosmological term.

Calculations $a(t)$ from $t = 10^{-43}$c to $t = 10^{-10}$c maked with the (19), for another points we use numerical calculations (18).

Notice, that use of our model formula for the Λ - term is unjustified, when we are speaking about the planck time. Therefore, values of the cosmological term and dark matter's densities in this times don't have pretensions of irrefragable coincidences with anticipated values. This results is used only for illustration of abrupt slump Λ-term in an Universe's first seconds. Nevertheless limited hard model of a dark matter assures the more steep inflation stage. Moreover we have for the planck time good coincidence with a planck matter density and expected value Λ-term $\Lambda \sim 10^{63} \dfrac{1}{\mathrm{sm}^2}$ until the phase transformation $SU(5) \longrightarrow SU(3)\times SU(2)\times U(1)$, which happened at $t = 10^{-35}$s and $T = 10^{15}$Gev . This features makes this model enough attractive. Notice, that in this model cosmological term is negative until $t \cong 5 \cdot 10^{16}$s $(a(t) \cong 5.75 \cdot 10^{27}\mathrm{sm})$. Anyway this model expects an additional researches. There notice, that limited hard state of a matter is complying with a system of particles with spin 1. The QFT's consideration of this system is showing a many interesting features [16]. Particularly, the particles in this model repels, but antiparticles gravitates. The particles which have a spin 1 is gluons and intermediate vector bosons. However, this particles are constrained and, consequently, haven't required for the limited hard model of a dark matter features.

Table 1. *Barion model of a dark model,* $k = 0.03$, $\Omega_0 = 0.6$

t (s)	$a(t)$, sm	$\Lambda = \Lambda_0 - kR$, sm^{-2}	ρ_D , $\frac{\text{g}}{\text{sm}^3}$
10^{-43}	$9 \cdot 10^{-4}$	$4 \cdot 10^{32}$	$2 \cdot 10^{60}$
10^{-35}	9	$9 \cdot 10^{20}$	$5 \cdot 10^{48}$
10^{-10}	$3 \cdot 10^{13}$	$4 \cdot 10^{-16}$	$2 \cdot 10^{12}$
10^{-5}	$8 \cdot 10^{15}$	$4 \cdot 10^{-23}$	$2 \cdot 10^{5}$
10^{-2}	$2 \cdot 10^{17}$	$4 \cdot 10^{-27}$	2
10^{-1}	$5 \cdot 10^{17}$	$3 \cdot 10^{-28}$	1
10^{0}	$2 \cdot 10^{18}$	$4 \cdot 10^{-30}$	$2 \cdot 10^{-2}$
10^{2}	$2 \cdot 10^{19}$	$6 \cdot 10^{-33}$	$3 \cdot 10^{-5}$
10^{7}	$6 \cdot 10^{21}$	$4 \cdot 10^{-40}$	$2 \cdot 10^{-12}$
10^{12}	$2 \cdot 10^{24}$	$2 \cdot 10^{-47}$	10^{-19}
10^{15}	10^{26}	$2 \cdot 10^{-52}$	10^{-24}
10^{16}	$6 \cdot 10^{26}$	10^{-54}	$6 \cdot 10^{-27}$
10^{17}	$2 \cdot 10^{27}$	$6 \cdot 10^{-56}$	$2 \cdot 10^{-28}$
10^{18}	$9 \cdot 10^{27}$	$2 \cdot 10^{-56}$	$2 \cdot 10^{-30}$

Table 2. *Limited hard model of a dark matter,* $k = 0.03$, $\Omega_0 = 0.63$

t (s)	$a(t)$, sm	$\Lambda = \Lambda_0 - kR$, sm^{-2}	ρ_D , $\frac{\text{g}}{\text{sm}^3}$
10^{-43}	$2 \cdot 10^9$	$-2 \cdot 10^{63}$	$5 \cdot 10^{90}$
10^{-35}	$7 \cdot 10^{11}$	$-6 \cdot 10^{46}$	$2 \cdot 10^{74}$
10^{-10}	$4 \cdot 10^{19}$	$-7 \cdot 10^{-4}$	$2 \cdot 10^{24}$
10^{-5}	10^{21}	$-7 \cdot 10^{-13}$	$2 \cdot 10^{15}$
10^{-2}	$9 \cdot 10^{21}$	$-5 \cdot 10^{-19}$	$2 \cdot 10^{9}$
10^{-1}	$2 \cdot 10^{22}$	$-3 \cdot 10^{-21}$	$9 \cdot 10^{6}$
10^{0}	$4 \cdot 10^{22}$	$-4 \cdot 10^{-23}$	10^{5}
10^{2}	$2 \cdot 10^{23}$	-10^{-27}	3
10^{7}	$6 \cdot 10^{24}$	$-3 \cdot 10^{-37}$	10^{-9}
10^{12}	$2 \cdot 10^{26}$	$-5 \cdot 10^{-47}$	$2 \cdot 10^{-19}$
10^{15}	$2 \cdot 10^{27}$	$-2 \cdot 10^{-53}$	$6 \cdot 10^{-26}$
10^{16}	$4 \cdot 10^{27}$	$-2 \cdot 10^{-55}$	$7 \cdot 10^{-28}$
10^{17}	$7 \cdot 10^{27}$	$2 \cdot 10^{-56}$	$2 \cdot 10^{-29}$
10^{18}	10^{28}	$2 \cdot 10^{-56}$	$2 \cdot 10^{-30}$

In conclusion we are going to consider the newtonian limit for our model. From [8] we have:

$$R_0^0 = \frac{1}{c^2}\Delta\varphi \ , T_0^0 = T = \varepsilon = \rho c^2 \ , \tag{20}$$

where φ is the gravity potential. As result, we take this equation:

$$\Delta\varphi = 4\pi G \frac{1-6k}{1-4k}\rho - \frac{c^2\Lambda_0}{1-4k} \ . \tag{21}$$

The item, which is proportional to Λ_0, evidently, incidentally for the newtonian limit. There we have, however, an important result: renormalization of the gravitational constant. It, as coefficient in the action for the gravitational field must be positive. From this we find one more condition $k < \frac{1}{6}$ or $k > \frac{1}{4}$. Our value $k = 0.03$ belongs to this interval and, moreover, in view of the fact his smallness, makes the difference between cosmological and newtonian values G very small : $G_{\rm Newt} \approx 0.93G$. This difference is incidentally too, because another cosmological parameters have larger errors, for example, Hubble's constant (11). Therefore we don't make a difference between renormalized and notrenormalized values of G.

3 Short conclusions

1). A model of a closed universe with a time-variant vacuum energy density is proposed.

2). It was shown that the presence of a variable $\Lambda-$term alters the evolution of the scale factor a in the very early epoch.

3). In the context of this model, a severe decrease of the vacuum energy density at the primitive stage of Universe's development is explained. The numerical results for the value of the Λ-term agree with theoretical predictions for the early epoch and with recent observation data.

4) Such explanations are speaking about more attractive the limited hard model of a dark matter, that requires additional its in-depth study.

5) As is shown, variable cosmological term is reducing to renormalization of the gravitational constant in the newtonian limit.

References

[1] Permulter S.J. et al. (1998) Nature **391** 51.

[2] Riess A.G. et al., Astron.J., **116**, (1998), 565.

[3] Perlmutter S. et al., Astrophys.J, **517**, (1999) 565.

[4] Kofman L.A., Gnedin N.Y., Bahcall N.A. (1993) Ap.J. **413**,1.

[5] Ostriker J.P. and Steinhardt P.J. (1995), Nature **377** 600.

[6] Weinberg C. Rev.Mod.Phys. (1989) v.61, p.1-23.

[7] Sahni V., Starobinsky A. arXiv:astro-ph/9904398 v.2 (2000).

[8] Landau L., Lifshic E. The Classical Theory of Fields (1951) (Pergamon, Oxford).

[9] Itzykson C. and Zuber J.-B. Quantum Field Theory. McGraw-Hill Book Company, 1082.

[10] Starobinskyy V. JETP Letters **30** 719 (1970).

[11] Birell N.D., Davies P.C.W. Quantum Fields in Curved Space. Cambridge University Press, 1982.

[12] Fomin P.I.Zero cosmological constant and Planck scales phenomenology. Proc. 4th seminar on quantum gravity (May,1987, Moscow). Ed. by M.markov.-Singapoure:World Scientific, 1988.-P.813.

[13] Sakharov A.D., JETP Letters,**5**, 1967, p. 32, 36 (in Russian)

[14] Chernin A.D. Usp.Fiz.Nauk v.171, № 11, (2001), p.1153.

[15] Peebles P.J.E. Principles of Physical Cosmology (Priceton, N.J. Princeton Univ.Press), 1993.

[16] Z'eldovich Ya. B., Novikov I.D., The theory of gravitation and star's evolution, Moscow, Nauka, 1971 (in Russian)